







COMPOSITION

Each Kilogram of **Bioshell** $^{\circ}$ Feed Premix contains 69.6 mg/Kg of 25 hydroxyvitamin D $_{3}$ and minimum 3.2x10 12 viable spores of Bacillus licheniformis

BENEFITS

versus vitamin D

- Bioshell® gives better zoo technical performance
- Bioshell® is more potent (young animals, mycotoxines, fatty liver...)
- Bioshell® establishes and maintains a beneficial microbial population in the gut by:
- Creating a favourable environment for beneficial bacteria via competitive exclusion
- Direct antagonism against Clostridium perfringens.

versus synthetic 25-OH D₃

- Bioshell® is 2 times more bio-available
- Bioshell® is more stable during storage and pelleting

versus $1,25-(OH)_2$ D₃ or 1α -OH D₃

- Bioshell® is absorbed in the egg yolk
- Bioshell[®] is safer in dosing

INDICATION

Bioshell[®] Feed Premix is used as an additive to compound feed of poultry (broilers, laying hens and poultry breeders) as a vitamin D₃ supplement to aid in maintaining calcium homeostasis and general bone health and Probiotic for improving gut health and better nutrient absorption.

DIRECTIONS FOR USE

Use as recommended by the nutritionist.

ROUTE OF ADMINISTRATION AND DOSAGE

Bioshell[®] should be added at the level of 250-500 gm per MT of finished feed.

WITHDRAWAL PERIOD

Zero days

PACK SIZE

Available in 1 Kg & 25 Kg

